



2. Graphical Transformations of Functions

In this section we will discuss how the graph of a function may be transformed either by shifting, stretching or compressing, or reflection. In this section let c be a positive real number.



Vertical Translations

A shift may be referred to as a translation. If c is added to the function, where the function becomes $y = f(x) + c$, then the graph of $f(x)$ will vertically shift upward by c units. If c is subtracted from the function, where the function becomes $y = f(x) - c$, then the graph of $f(x)$ will vertically shift downward by c units. In general, a vertical translation means that every point (x, y) on the graph of $f(x)$ is transformed to $(x, y + c)$ or $(x, y - c)$ on the graphs of $y = f(x) + c$ or $y = f(x) - c$ respectively.





Horizontal Translations

If c is added to the variable of the function, where the function becomes $y = f(x + c)$, then the graph of $f(x)$ will horizontally shift to the left c units. If c is subtracted from the variable of the function, where the function becomes $y = f(x - c)$, then the graph of $f(x)$ will horizontally shift to the right c units. In general, a horizontal translation means that every point (x, y) on the graph of $f(x)$ is transformed to $(x - c, y)$ or $(x + c, y)$ on the graphs of $y = f(x + c)$ or $y = f(x - c)$ respectively.





Reflection

If the function or the variable of the function is multiplied by -1, the graph of the function will undergo a reflection. When the function is multiplied by -1 where $y = f(x)$ becomes $y = -f(x)$, the graph of $y = f(x)$ is reflected across the x-axis.




On the other hand, if the variable is multiplied by -1, where $y = f(x)$ becomes $y = f(-x)$, the graph of $y = f(x)$ is reflected across the y-axis.




Vertical Stretching and Shrinking

If c is multiplied to the function then the graph of the function will undergo a vertical stretching or compression. So when the function becomes $y = cf(x)$ and $0 < c < 1$, a vertical shrinking of the graph of $y = f(x)$ will occur. Graphically, a vertical shrinking pulls the graph of $y = f(x)$ toward the x -axis. When $c > 1$ in the function $y = cf(x)$, a vertical stretching of the graph of $y = f(x)$ will occur. A vertical stretching pushes the graph of $y = f(x)$ away from the x -axis. In general, a vertical stretching or shrinking means that every point (x, y) on the graph of $f(x)$ is transformed to (x, cy) on the graph of $y = cf(x)$.



Horizontal Stretching and Shrinking

If c is multiplied to the variable of the function then the graph of the function will undergo a horizontal stretching or compression. So when the function becomes $y = f(cx)$ and $0 < c < 1$, a horizontal stretching of the graph of $y = f(x)$ will occur. Graphically, a horizontal stretching pulls the graph of $y = f(x)$ away from the y -axis. When $c > 1$ in the function $y = f(cx)$, a horizontal shrinking of the graph of $y = f(x)$ will occur. A horizontal shrinking pushes the graph of $y = f(x)$ toward the y -axis. In general, a horizontal stretching or shrinking means that every point (x, y) on the graph of $f(x)$ is transformed to $(x/c, y)$ on the graph of $y = f(cx)$.



Transformations can be combined within the same function so that one graph can be shifted, stretched, and reflected. If a function contains more than one transformation it may be graphed using the following procedure:


Steps for Multiple Transformations

Use the following order to graph a function involving more than one transformation:

1. Horizontal Translation
2. Stretching or shrinking
3. Reflecting
4. Vertical Translation


Examples: Graph the following functions and state their domain and range:

1. $f(x) = (x + 2)^2 - 3$
basic function (b.f.) = x^2 , $\leftarrow 2, \downarrow 3$



$$\text{Domain} = (-\infty, \infty)$$
$$\text{Range} = [-3, \infty)$$


2. $f(x) = -|x - 3| + 1$
 b.f. = $|x|$, $\rightarrow 3$, reflect about x-axis, $\uparrow 1$



Domain = $(-\infty, \infty)$
 Range = $(-\infty, 1]$


3. $f(x) = -2\sqrt{x+3} + 1$

b.f. = \sqrt{x} , $\leftarrow 3$, stretch about y-axis, reflect about x-axis, $\uparrow 1$



Domain = $[-3, \infty)$
 Range = $(-\infty, 1]$


4. $f(x) = (-2x + 1)^3 - 2$
 b.f. = x^3 , $\leftarrow 1$, shrink about x-axis ($c = 2$), reflect about y-axis, $\downarrow 2$



Domain = $(-\infty, \infty)$


Range = $(-\infty, \infty)$

5. Let the graph of $f(x)$ be the following:




Graph the following problems:


a. $f(x) - 3$




b. $f(x + 1)$




c. $f(x - 2) + 1$



d. $-f(x + 1)$



e. $f(-x) - 2$



Transformations of the graphs of functions

$f(x) + c$	shift $f(x)$ up c units
$f(x) - c$	shift $f(x)$ down c units
$f(x + c)$	shift $f(x)$ left c units
$f(x - c)$	shift $f(x)$ right c units
$f(-x)$	reflect $f(x)$ about the y-axis
$-f(x)$	reflect $f(x)$ about the x-axis
$cf(x)$	When $0 < c < 1$ – vertical shrinking of $f(x)$ When $c > 1$ – vertical stretching of $f(x)$ Multiply the y values by c
$f(cx)$	When $0 < c < 1$ – horizontal stretching of $f(x)$ When $c > 1$ – horizontal shrinking of $f(x)$ Divide the x values by c